

**HEREDITARY ANGIOEDEMA (HAE) ENROLLMENT FORM**

**1 PATIENT INFORMATION**  
*(Please complete the following or send patient demographic sheet)*

Patient Name: \_\_\_\_\_  
 Primary Phone: \_\_\_\_\_  
 DOB: \_\_\_\_\_ Gender: \_\_\_\_\_  
 Address: \_\_\_\_\_  
 \_\_\_\_\_  
 Allergy: \_\_\_\_\_  
 Patient Weight: \_\_\_\_\_  
 Diagnosis: D84.1 Other \_\_\_\_\_

**2 PRESCRIBER INFORMATION**

Prescriber's Name: \_\_\_\_\_  
 Contact Person: \_\_\_\_\_ Phone: \_\_\_\_\_  
 NPI #: \_\_\_\_\_  
 Address: \_\_\_\_\_  
 \_\_\_\_\_  
 City, State, Zip: \_\_\_\_\_  
 Phone: \_\_\_\_\_ Fax: \_\_\_\_\_

**3 INSURANCE INFORMATION** *(Please copy and attach the front and back of insurance and prescription drug card)*

**4 CATHETER ACCESS AND FLUSH PROTOCOL** (If Applicable)

ACCESS TYPE	CATHETER FLUSH ORDERS
Peripheral	<b>0.9% Saline Flush: Dispense: 30 Days Refills: PRN x 1yr</b> Flush line/port with 10mL for patency/SASH protocol.
PORT <i>(Also include Peripheral IV PRN Port Malfunction)</i>	<b>Heparin Flush: Dispense: 30 Days Refills: PRN x 1yr</b> Flush port with _____ mL of Heparin _____ units/mL per SASH protocol.

**5 PRESCRIPTION INFORMATION** Please check the following:

MEDICATION	DOSE	DIRECTIONS	DAY SUPPLY	QUANTITY	REFILLS
<input type="checkbox"/> BERINERT® (C1 Esterase inhibitor [human])	20 IU/kg	_____	_____	_____	_____
<input type="checkbox"/> CINRYZE® (C1 esterase inhibitor [human])	max 100 IU/kg up to 2500 IU	_____	_____	_____	_____
<input type="checkbox"/> FIRAZYR® (Icatibant injection)	30 mg prefilled syringe	Inject 1 syringe (30mg) subcutaneously in the abdominal area. If response is inadequate or symptoms recur, additional injections of 30mg may be administered at intervals of at least 6 hours.	_____	_____	_____
<input type="checkbox"/> HAEGARDA® C1 Esterase Inhibitor Subcutaneous [human]	max 60 IU/kg	_____	28 Days	<input type="checkbox"/> 2000 IU #of vials _____ <input type="checkbox"/> 3000 IU #of vials _____	13
<input type="checkbox"/> KALBITOR® (ecallantide) injection	30 mg	Administer 30mg (3mL) SC in three 10mg (1mL) injections as needed for acute HAE attack. Dose may be repeated within a 24 hour period.	_____	_____ boxes of three 10mg (1mL) vials	_____
<input type="checkbox"/> RUCONEST® (C1 esterase inhibitor [recombinant])	50 IU/kg max. 4200 units	_____	_____	_____	_____
<input type="checkbox"/> TAKHZYRO® (Ivanide) injection	150 mg 300 mg	Administer subcutaneously every 2 weeks (bi-weekly) Administer subcutaneously once monthly (monthly)	28 Days	2 pre-filled syringes (biweekly) 1 pre-filled syringe (monthly)	13

**6 PHYSICIAN SIGNATURE** (Required)

X \_\_\_\_\_ Date of Signature  
 PRODUCT SUBSTITUTION PERMITTED

X \_\_\_\_\_ Date of Signature  
 DISPENSE AS WRITTEN

IMPORTANT NOTICE: This facsimile transmission is intended to be delivered only to the named addressee and may contain material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If it is received by anyone other than the named addressee, the recipient should immediately notify the sender at the address and telephone number set forth herein and obtain instructions as to disposal of the transmitted material. In no event should such material be read or retained by anyone other than the named addressee, except by express authority of the sender to the named addressee.

**HEREDITARY ANGIOEDEMA (HAE) PRESCRIPTION FORM**

**7 ADDITIONAL INSTRUCTIONS**

**SITE OF CARE**

- Self/caregiver administration training # visits ordered \_\_\_\_\_ or  competent
- Home Health Nursing

**NURSING INSTRUCTIONS**

1. Gain IV access prior to mixing (if applicable).
2. Mix and administer \_\_\_\_\_ according to package insert (main recommendation)

**IN CASE OF EMERGENCY**

1. Stop medication
2. Call doctor
3. Administer emergency med if ordered in box \_\_\_\_\_

**ADMINISTER EMERGENCY MEDS PER PHYSICIAN ORDERS** #q.s. for each drug. Refill: PRN x 1 year

For severe anaphylaxis, administer prescribed epinephrine.

If severe symptoms persist, may repeat. **(Please select epinephrine dose):**

epinephrine 0.3 mg autoinjector IM (patients >30 kg)

epinephrine 0.15 mg autoinjector IM (patients <30 kg)

- Diphenhydramine \_\_\_\_\_ mg IV push over 2-5 minutes for Infusion Reaction
- Corticosteroid (specify drug and dose):

Other: \_\_\_\_\_

**For severe hypersensitive reaction, stop infusion, administer epinephrine Autoinjector IM - may repeat in 20 minutes if needed. Call 911.**

**8 PHYSICIAN SIGNATURE (Required)**

X \_\_\_\_\_ X \_\_\_\_\_  
 PRODUCT SUBSTITUTION PERMITTED Date of Signature DISPENSE AS WRITTEN Date of Signature

IMPORTANT NOTICE: This facsimile transmission is intended to be delivered only to the named addressee and may contain material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If it is received by anyone other than the named addressee, the recipient should immediately notify the sender at the address and telephone number set forth herein and obtain instructions as to disposal of the transmitted material. In no event should such material be read or retained by anyone other than the named addressee, except by express authority of the sender to the named addressee.